9142使用书-2闭
料号：2-0023-2410-000
尺寸：40x140mm
材质：70g书写纸
颜色：黑白印刷
折页

EN

DISCLAIMER
Please read this user manual before use. Failure to follow the instructions for proper setup, use and care for your device can increase the risk of serious personal injury, death, or property damage.

The manufacturer assumes no responsibility for any damage or injury caused by improper handling or beyond normal usage without it's intended.

RECOMMENDED SETUP
We recommend the exclusive use of Aspire original components and accessories. Using non-Aspire replacement or third-party parts may damage the device and cause overheating or excessive heat conditions (167°C / 154°F).

Your device can be damaged if dropped, burned, punctured, crushed, or if it comes in contact with liquid. If you suspect damage to your device or its battery, discontinue use immediately. Damaged devices can overheating or burn.

To avoid malfunctions, always ensure you either lock or switch off your device when not in use, especially when carrying your device in your pocket, bag, or purse.

In the case of burn or abnormal deformation of one of the components, please stop using immediately and contact Aspire Customer Care.

COIL MAINTENANCE SAFETY
A coil that not functioning or cold to burn the cotton dry hot shots at empty puff. Let the puffers stand for 5 minutes after filling so the cotton can be saturated with e-liquid before use.

REPLACEABLE BATTERIES SAFETY
Do not use non-rechargeable batteries. Do not use damaged batteries or batteries with a torn or damaged wrapper. Disconnect the battery from the charger when it is not in use. Do not use your vape device while charging.

PRODUCT DISPOSAL SAFETY
Do not discard this product and battery in the environment. Please dispose of the product components or accessories in collection/disposal points.

HEALTH WARNING SAFETY
Do not use this product if you are pregnant, breastfeeding, generally ill, or suffer from respiratory and/or cardiovascular system disorders.

If you experience any side effects discontinue using the product. If the effect(s) continue after discontinuing, consult a healthcare professional immediately.

Package Contents:
* Huracan EX Mod
* 1* Huracan Tank (2ml)
* 1* Replacement Glass (2ml) (pre-installed)
* 1* Spare Bubble Glass Tube
* 1* Spare O-Ring
* 1* USB-C Cable
* User Manual

Components View:

Kit Dimensions: 136.75x132.5x mm

Tank Specifications:
810 Replaceable drip tip
Tank Capacity: 6.5 ml
Tank Diameter: 25.4 mm
Top Fill Port - Rubber Stopper
Coil Compatibility: Replaceable BF resistant coils compatible
Tank Material: PC, Pyrex, Steel
Tank Weight: 127.5 g
Tank Dimensions (Including 810 Drip Tip): 67.5x41.5x mm

Mod Specifications:
Mod Material: Zinc Alloy
BF resistant Coil: 0.4Ω (0.28-0.35Ω)
Drip Tip: 810 (15.25x8)
Mod Weight: 127.5 g
Dimensions (Including 810 Drip Tip): 67.5x41.5x mm

Coil Installation:
1. Unscrew to remove the bottom hardware with Horus.
2. Put the pre-used coil with more force and push the new coil into the tank.
3. Screw the bottom hardware back with force.



Airpressure Setting:
Step 1:
Enter the system interface, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS".
Step 2:
Press the "Fire" button continuously to enter the menu adjustment.

Instructions:

Mod Condition	Status of Indicator Light & Details	
Turn on / off	Toggle the power switch to turn the device on / off.	Normal Operation
Lock or unlock the mod	Toggle the power switch to lock / unlock the device.	Normal Operation
Lock or unlock the mod	Press the "+" & "-" buttons simultaneously to lock / unlock the device. Lock the buttons prevent only the Firebutton and not the Firebutton.	Normal Operation
Screen brightness adjustment	Press the button "Video" (Horus) or "Fire" button (Aspire) twice. Only the Horus Firebutton can adjust the screen brightness.	Normal Operation
Enter the menu	Press the button "Video" (Horus) or "Fire" button (Aspire) continuously. Release the button when the screen starts. Only the Horus Firebutton can enter the menu.	Normal Operation
Adjust the Airpressure	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to adjust the airpressure.	Normal Operation
Adjust the Wattage	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to adjust the wattage.	Normal Operation
Adjust the Voltage	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to adjust the voltage.	Normal Operation
Adjust the Puffs	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to adjust the puffs.	Normal Operation
Adjust the Airpressure	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to adjust the airpressure.	Normal Operation
Reset the Puffs	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to reset the puffs.	Normal Operation
Reset the Puffs	Enter the menu, the screen will display "BATTERY", "VOLT", "WATTAGE", "VOLTAGE", "CURRENT", "RESIST OHOS", and "PUFFS". Press the button "+" or "-" to reset the puffs.	Normal Operation
Remember to close the combination in the tank.		

Remember to close the combination in the tank.

Note:
* When installing a new coil, let the tank stand for 5 minutes to ensure the cotton is fully saturated with e-liquid for optimal flavor.
* Regularly check the cotton for any damage. Refill the tank to avoid dry burning or damage to the coil.

Safety Features:

Vaping Time Cap (Protection Mode)	The system will slightly reduce the power.
Clean Circuit Protection	The system will shut down.
Short Circuit Protection	The system will shut down.
Low Resistance Protection	The screen will display "LOW SOC".
Low Voltage Protection	The screen will display "LOW SOC".
Overcharge Protection	The screen will display "FULL".
Over Heat Protection	The screen will display "TOO HOT".
Abnormal Charge Protection	The screen will display "CHECK USB".
Battery Reverse Protection	The screen will stop working.

Battery Installation:



Charging:

Maximum charge current	2A
Maximum charge voltage	5V
Fully charged	The screen will display no pattern.
Fully charged	The screen will display "FULL".
Fully charged (Lock the device in lock mode)	The screen will display "CHECK USB" for 3 seconds.
Fully charged (Lock the device in lock mode)	The screen will stop working. Reliability will stop charging to ensure safety.

IMPORTANT INFORMATION ABOUT USING NICOTINE CONTAINING E-LIQUIDS

1. NICOTINE DEPENDENCE & LIQUIDS
Before using nicotine containing e-liquid for sale, should read the following information carefully.

2. NICOTINE CONTAINING E-LIQUID
If used with nicotine containing liquid, it should be noted that nicotine is a highly toxic substance. Even in small quantities, it can be absorbed by the skin and cause toxic effects if consumed orally, especially in infants or children.

Do not use if you are allergic to nicotine or any of the other ingredients in the e-liquid formula.

Nicotine can be absorbed through the skin.

3. NICOTINE WARNINGS AND PRECAUTIONS
Please carefully read the following information:
* Non-smokers
* Do not use if you are under the legal age.
* Nicotine containing e-liquid and the nicotine is known to cause the addiction and to be excreted in breast milk.

4. EFFECTS DESCRIBED
* These experiencing general ill health, such as heart problems, lung problems, diabetes, ulcers, or high blood pressure, should consult their doctor before use, as these conditions may be exacerbated by nicotine.
* Nicotine may affect the blood circulation and have a pharmacological effect on some of the vascular beds that can affect high blood pressure.
* Nicotine may affect the digestive system, including bowel and bladder.
* These taking the following medication:
* Insulin, Nicotine may affect blood sugar levels.
* Rizamatan (to treat restless leg syndrome)
* Tricyclene (to treat schizophrenia)

5. POSSIBLE SIDE EFFECTS
If you inhale too much nicotine or use the product too much, you may experience one or more of the following effects:
* Feeling faint
* Nausea (feeling sick)
* Headache
* Coughing
* Irritation of the mouth or throat
* Stomach pain
* Diarrhea
* Hiccups
* Palpitations
* Vomiting
* Arrhythmia in the chest

6. POSSIBLE LONG TERM EFFECTS
Regular use of nicotine containing e-liquid may cause the following:
* Nausea, Vomiting
* Headache
* Dizziness
* Tiredness
* Tremors
* Cardiac Arrhythmia
* Chest Pain
* Hypertension
* Coronary Artery Disease
* Stroke
* Cancer
* Nasal Congestion
* Sinusitis
* Allergies
* Bronchitis
* Emphysema
* Gastrointestinal Disorders
* Peptic Ulcer
* Gastritis
* Diarrhea
* Constipation
* Nausea
* Vomiting
* Gastroesophageal Reflux Disease (GERD)
* Erectile Dysfunction
* Premature Ejaculation
* Reduced Libido
* Menstrual Irregularities
* Fertility Issues
* Pregnancy Complications
* Low Birth Weight
* Premature Birth
* Stillbirth
* Sudden Infant Death Syndrome (SIDS)
* Impotence
* Osteoporosis
* Muscle Weakness
* Joint Pain
* Rheumatoid Arthritis
* Psoriasis
* Eczema
* Premature Aging
* Wrinkles
* Yellowing of Teeth
* Bad Breath
* Gum Disease
* Tooth Decay
* Tooth Loss
* Cognitive Decline
* Memory Loss
* Depression
* Anxiety
* Insomnia
* Sleep Disorders
* Fatigue
* Weakness
* Muscle Cramps
* Numbness
* Tingling
* Chronic Pain
* Headaches
* Migraines
* Vertigo
* Tinnitus
* Hearing Loss
* Vision Problems
* Cataracts
* Glaucoma
* Macular Degeneration
* Dry Eyes
* Conjunctivitis
* Eye Strain
* Blurred Vision
* Dizziness
* Balance Problems
* Coordination Issues
* Memory Problems
* Confusion
* Mood Swings
* Irritability
* Aggressive Behavior
* Addiction
* Withdrawal Symptoms
* Cravings
* Relapse
* Social Isolation
* Financial Problems
* Legal Issues
* Family Conflicts
* Loss of Productivity
* Decreased Performance
* Absenteeism
* Unemployment
* Homelessness
* Poverty
* Malnutrition
* Dehydration
* Hypothermia
* Frostbite
* Gangrene
* Amputation
* Death

7. CONTRAINDICATIONS
* Do not use if you are allergic to the preferred general contents of the formulation (propylene glycol, glycerine and nicotine) or products/materials (e.g. rubbers or plastics, components or accessories) that come into contact with the e-liquid.
* Do not use if you are under the legal age.
* Do not use if you are pregnant or breastfeeding.
* Do not use if you are in the process of quitting nicotine.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.

8. PROBLEMS EFFECTS SECONDARIES
* Nicotine containing products or other products or services provided complement your consumption of nicotine cigarettes effects:
* Dehydration
* Nausea (sensation of nausea)
* Dizziness
* Headache
* Irritation of the throat or gargling
* Malaise
* Vertigo
* Cough
* Congestion nasal
* Cerebral
* Allergies
* Cardiac palpitations

9. CONTRAINDICATIONS
* Do not use if you are allergic to the preferred general contents of the formulation (propylene glycol, glycerine and nicotine) or products/materials (e.g. rubbers or plastics, components or accessories) that come into contact with the e-liquid.
* Do not use if you are under the legal age.
* Do not use if you are pregnant or breastfeeding.
* Do not use if you are in the process of quitting nicotine.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.

10. ADVERTENCIAS Y PRECAUCIONES SOBRE LA NICOTINA
Antes de utilizar el e-líquido que contiene nicotina, es aquello que debe tener en cuenta:
* Fumadores
* Prohibido a menores de edad.
* Nicotina es conocida por ser adictiva y se excreta en la leche materna.

11. EFECTOS DESCRITOS
* Aquellos que tienen problemas de salud en general, como problemas del corazón, pulmones, diabetes, ulceras o presión arterial alta, deben consultar a su médico antes de usar, ya que estas condiciones pueden ser exacerbadas por la nicotina.
* La nicotina puede afectar la circulación sanguínea y tener un efecto farmacológico en algunos lechos vasculares que pueden afectar la presión arterial.
* La nicotina puede afectar el sistema digestivo, incluyendo intestinos y vejiga.
* Aquellos que toman:
* Insulina, la nicotina puede afectar los niveles de azúcar en sangre.
* Rizamatan (para tratar el síndrome de las piernas inquietas)
* Triciclene (para tratar la esquizofrenia)

12. PROBLEMAS EFECTOS SECUNDARIOS
Si inhala demasiada nicotina o usa el producto demasiado, puede experimentar uno o más de los siguientes efectos:
* Mareos
* Náuseas (sensación de mareo)
* Dolor de cabeza
* Tos
* Irritación de la boca o la garganta
* Dolor de estómago
* Diarrea
* Hipo
* Palpitaciones
* Vomitar
* Arritmia en el pecho

13. PROBLEMAS EFECTOS SECUNDARIOS
Si se usa nicotina con regularidad, puede provocar:
* Náuseas, vomitar
* Dolor de cabeza
* Mareos
* Fatiga
* Temblores
* Arritmia cardíaca
* Dolor en el pecho
* Hipertensión
* Enfermedad arterial coronaria
* Accidente cerebrovascular
* Cáncer
* Congestión nasal
* Sinusitis
* Alergias
* Bronquitis
* Enfisema
* Trastornos gastrointestinales
* Úlcera péptica
* Gastritis
* Diarrea
* Estreñimiento
* Náuseas
* Vomitar
* Enfermedad por reflujo gastroesofágico (ERGE)
* Disfunción eréctil
* Eyaculación precoz
* Disminución de la libido
* Irregularidades menstruales
* Problemas de fertilidad
* Complicaciones del embarazo
* Bajo peso al nacer
* Parto prematuro
* Mortinato
* Síndrome de muerte súbita del lactante (SMSL)
* Impotencia
* Osteoporosis
* Debilidad muscular
* Dolor articular
* Artritis reumatoide
* Psoriasis
* Eczema
* Envejecimiento prematuro
* Arrugas
* Amarillamiento de los dientes
* Mal aliento
* Enfermedades de las encías
* Caries dental
* Pérdida dental
* Deterioro cognitivo
* Pérdida de memoria
* Confusión
* Depresión
* Ansiedad
* Insomnio
* Trastornos del sueño
* Fatiga
* Debilidad
* Calambres musculares
* Entumecimiento
* Hormigueo
* Dolor crónico
* Dolores de cabeza
* Migrañas
* Vértigo
* Tinnitus
* Pérdida de audición
* Problemas de visión
* Cataratas
* Glaucoma
* Degeneración macular
* Ojos secos
* Conjuntivitis
* Fatiga ocular
* Visión borrosa
* Mareos
* Problemas de equilibrio
* Problemas de coordinación
* Problemas de memoria
* Confusión
* Cambios de humor
* Irritabilidad
* Comportamiento agresivo
* Adicción
* Síntomas de abstinencia
* Antojos
* Recaída
* Aislamiento social
* Problemas financieros
* Problemas legales
* Conflictos familiares
* Pérdida de productividad
* Disminución del rendimiento
* Ausentismo
* Desempleo
* Pobreza
* Malnutrición
* Deshidratación
* Hipotermia
* Congelación
* Gangrena
* Amputación
* Muerte

14. ADVERTENCIAS Y PRECAUCIONES SOBRE LA NICOTINA
Antes de utilizar el e-líquido que contiene nicotina, es aquello que debe tener en cuenta:
* Fumadores
* Prohibido a menores de edad.
* Nicotina es conocida por ser adictiva y se excreta en la leche materna.

15. EFECTOS DESCRITOS
* Aquellos que tienen problemas de salud en general, como problemas del corazón, pulmones, diabetes, ulceras o presión arterial alta, deben consultar a su médico antes de usar, ya que estas condiciones pueden ser exacerbadas por la nicotina.
* La nicotina puede afectar la circulación sanguínea y tener un efecto farmacológico en algunos lechos vasculares que pueden afectar la presión arterial.
* La nicotina puede afectar el sistema digestivo, incluyendo intestinos y vejiga.
* Aquellos que toman:
* Insulina, la nicotina puede afectar los niveles de azúcar en sangre.
* Rizamatan (para tratar el síndrome de las piernas inquietas)
* Triciclene (para tratar la esquizofrenia)

16. PROBLEMAS EFECTOS SECUNDARIOS
Si inhala demasiada nicotina o usa el producto demasiado, puede experimentar uno o más de los siguientes efectos:
* Mareos
* Náuseas (sensación de mareo)
* Dolor de cabeza
* Tos
* Irritación de la boca o la garganta
* Dolor de estómago
* Diarrea
* Hipo
* Palpitaciones
* Vomitar
* Arritmia en el pecho

17. PROBLEMAS EFECTOS SECUNDARIOS
Si se usa nicotina con regularidad, puede provocar:
* Náuseas, vomitar
* Dolor de cabeza
* Mareos
* Fatiga
* Temblores
* Arritmia cardíaca
* Dolor en el pecho
* Hipertensión
* Enfermedad arterial coronaria
* Accidente cerebrovascular
* Cáncer
* Congestión nasal
* Sinusitis
* Alergias
* Bronquitis
* Enfisema
* Trastornos gastrointestinales
* Úlcera péptica
* Gastritis
* Diarrea
* Estreñimiento
* Náuseas
* Vomitar
* Enfermedad por reflujo gastroesofágico (ERGE)
* Disfunción eréctil
* Eyaculación precoz
* Disminución de la libido
* Irregularidades menstruales
* Problemas de fertilidad
* Complicaciones del embarazo
* Bajo peso al nacer
* Parto prematuro
* Mortinato
* Síndrome de muerte súbita del lactante (SMSL)
* Impotencia
* Osteoporosis
* Debilidad muscular
* Dolor articular
* Artritis reumatoide
* Psoriasis
* Eczema
* Envejecimiento prematuro
* Arrugas
* Amarillamiento de los dientes
* Mal aliento
* Enfermedades de las encías
* Caries dental
* Pérdida dental
* Deterioro cognitivo
* Pérdida de memoria
* Confusión
* Depresión
* Ansiedad
* Insomnio
* Trastornos del sueño
* Fatiga
* Debilidad
* Calambres musculares
* Entumecimiento
* Hormigueo
* Dolor crónico
* Dolores de cabeza
* Migrañas
* Vértigo
* Tinnitus
* Pérdida de audición
* Problemas de visión
* Cataratas
* Glaucoma
* Degeneración macular
* Ojos secos
* Conjuntivitis
* Fatiga ocular
* Visión borrosa
* Mareos
* Problemas de equilibrio
* Problemas de coordinación
* Problemas de memoria
* Confusión
* Cambios de humor
* Irritabilidad
* Comportamiento agresivo
* Adicción
* Síntomas de abstinencia
* Antojos
* Recaída
* Aislamiento social
* Problemas financieros
* Problemas legales
* Conflictos familiares
* Pérdida de productividad
* Disminución del rendimiento
* Ausentismo
* Desempleo
* Pobreza
* Malnutrición
* Deshidratación
* Hipotermia
* Congelación
* Gangrena
* Amputación
* Muerte

IMPORTANT INFORMATION ABOUT USING NICOTINE CONTAINING E-LIQUIDS

1. NICOTINE DEPENDENCE & LIQUIDS
Before using nicotine containing e-liquid for sale, should read the following information carefully.

2. NICOTINE CONTAINING E-LIQUID
If used with nicotine containing liquid, it should be noted that nicotine is a highly toxic substance. Even in small quantities, it can be absorbed by the skin and cause toxic effects if consumed orally, especially in infants or children.

Do not use if you are allergic to nicotine or any of the other ingredients in the e-liquid formula.

Nicotine can be absorbed through the skin.

3. NICOTINE WARNINGS AND PRECAUTIONS
Please carefully read the following information:
* Non-smokers
* Do not use if you are under the legal age.
* Nicotine containing e-liquid and the nicotine is known to cause the addiction and to be excreted in breast milk.

4. EFFECTS DESCRIBED
* These experiencing general ill health, such as heart problems, lung problems, diabetes, ulcers, or high blood pressure, should consult their doctor before use, as these conditions may be exacerbated by nicotine.
* Nicotine may affect the blood circulation and have a pharmacological effect on some of the vascular beds that can affect high blood pressure.
* Nicotine may affect the digestive system, including bowel and bladder.
* These taking the following medication:
* Insulin, Nicotine may affect blood sugar levels.
* Rizamatan (to treat restless leg syndrome)
* Tricyclene (to treat schizophrenia)

5. POSSIBLE SIDE EFFECTS
If you inhale too much nicotine or use the product too much, you may experience one or more of the following effects:
* Feeling faint
* Nausea (feeling sick)
* Headache
* Coughing
* Irritation of the mouth or throat
* Stomach pain
* Diarrhea
* Hiccups
* Palpitations
* Vomiting
* Arrhythmia in the chest

6. POSSIBLE LONG TERM EFFECTS
Regular use of nicotine containing e-liquid may cause the following:
* Nausea, Vomiting
* Headache
* Dizziness
* Tiredness
* Tremors
* Cardiac Arrhythmia
* Chest Pain
* Hypertension
* Coronary Artery Disease
* Stroke
* Cancer
* Nasal Congestion
* Sinusitis
* Allergies
* Bronchitis
* Emphysema
* Gastrointestinal Disorders
* Peptic Ulcer
* Gastritis
* Diarrhea
* Constipation
* Nausea
* Vomiting
* Gastroesophageal Reflux Disease (GERD)
* Erectile Dysfunction
* Premature Ejaculation
* Reduced Libido
* Menstrual Irregularities
* Fertility Issues
* Pregnancy Complications
* Low Birth Weight
* Premature Birth
* Stillbirth
* Sudden Infant Death Syndrome (SIDS)
* Impotence
* Osteoporosis
* Muscle Weakness
* Joint Pain
* Rheumatoid Arthritis
* Psoriasis
* Eczema
* Premature Aging
* Wrinkles
* Yellowing of Teeth
* Bad Breath
* Gum Disease
* Tooth Decay
* Tooth Loss
* Cognitive Decline
* Memory Loss
* Depression
* Anxiety
* Insomnia
* Sleep Disorders
* Fatigue
* Weakness
* Muscle Cramps
* Numbness
* Tingling
* Chronic Pain
* Headaches
* Migraines
* Vertigo
* Tinnitus
* Hearing Loss
* Vision Problems
* Cataracts
* Glaucoma
* Macular Degeneration
* Dry Eyes
* Conjunctivitis
* Eye Strain
* Blurred Vision
* Dizziness
* Balance Problems
* Coordination Issues
* Memory Problems
* Confusion
* Mood Swings
* Irritability
* Aggressive Behavior
* Addiction
* Withdrawal Symptoms
* Cravings
* Relapse
* Social Isolation
* Financial Problems
* Legal Issues
* Family Conflicts
* Loss of Productivity
* Decreased Performance
* Absenteeism
* Unemployment
* Homelessness
* Poverty
* Malnutrition
* Dehydration
* Hypothermia
* Frostbite
* Gangrene
* Amputation
* Death

7. CONTRAINDICATIONS
* Do not use if you are allergic to the preferred general contents of the formulation (propylene glycol, glycerine and nicotine) or products/materials (e.g. rubbers or plastics, components or accessories) that come into contact with the e-liquid.
* Do not use if you are under the legal age.
* Do not use if you are pregnant or breastfeeding.
* Do not use if you are in the process of quitting nicotine.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.

8. PROBLEMS EFFECTS SECONDARIES
* Nicotine containing products or other products or services provided complement your consumption of nicotine cigarettes effects:
* Dehydration
* Nausea (sensation of nausea)
* Dizziness
* Headache
* Irritation of the throat or gargling
* Malaise
* Vertigo
* Cough
* Congestion nasal
* Cerebral
* Allergies
* Cardiac palpitations

9. CONTRAINDICATIONS
* Do not use if you are allergic to the preferred general contents of the formulation (propylene glycol, glycerine and nicotine) or products/materials (e.g. rubbers or plastics, components or accessories) that come into contact with the e-liquid.
* Do not use if you are under the legal age.
* Do not use if you are pregnant or breastfeeding.
* Do not use if you are in the process of quitting nicotine.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.
* Do not use if you are in the process of quitting nicotine containing products.

10. ADVERTENCIAS Y PRECAUCIONES SOBRE LA NICOTINA
Antes de utilizar el e-líquido que contiene nicotina, es aquello que debe tener en cuenta:
* Fumadores
* Prohibido a menores de edad.
* Nicotina es conocida por ser adictiva y se excreta en la leche materna.

11. EFECTOS DESCRITOS
* Aquellos que tienen problemas de salud en general, como problemas del corazón, pulmones, diabetes, ulceras o presión arterial alta, deben consultar a su médico antes de usar, ya que estas condiciones pueden ser exacerbadas por la nicotina.
* La nicotina puede afectar la circulación sanguínea y tener un efecto farmacológico en algunos lechos vasculares que pueden afectar la presión arterial.
* La nicotina puede afectar el sistema digestivo, incluyendo intestinos y vejiga.
* Aquellos que toman:
* Insulina, la nicotina puede afectar los niveles de azúcar en sangre.
* Rizamatan (para tratar el síndrome de las piernas inquietas)
* Triciclene (para tratar la esquizofrenia)

12. PROBLEMAS EFECTOS SECUNDARIOS
Si inhala demasiada nicotina o usa el producto demasiado, puede experimentar uno o más de los siguientes efectos:
* Mareos
* Náuseas (sensación de mareo)
* Dolor de cabeza
* Tos
* Irritación de la boca o la garganta
* Dolor de estómago
* Diarrea
* Hipo
* Palpitaciones
* Vomitar
* Arritmia en el pecho

13. PROBLEMAS EFECTOS SECUNDARIOS
Si se usa nicotina con regularidad, puede provocar:
* Náuseas, vomitar
* Dolor de cabeza
* Mareos
* Fatiga
* Temblores
* Arritmia cardíaca
* Dolor en el pecho
* Hipertensión
* Enfermedad arterial coronaria
* Accidente cerebrovascular
* Cáncer
* Congestión nasal
* Sinusitis
* Alergias
* Bronquitis
* Enfisema
* Trastornos gastrointestinales
* Úlcera péptica
* Gastritis
* Diarrea
* Estreñimiento
* Náuseas
* Vomitar
* Enfermedad por reflujo gastroesofágico (ERGE)
* Disfunción eréctil
* Eyaculación precoz
* Disminución de la libido
* Irregularidades menstruales
* Problemas de fertilidad
* Complicaciones del embarazo
* Bajo peso al nacer
* Parto prematuro
* Mortinato
* Síndrome de muerte súbita del lactante (SMSL)
* Impotencia
* Osteoporosis
* Debilidad muscular
* Dolor articular
* Artritis reumatoide
* Psoriasis
* Eczema
* Envejecimiento prematuro
* Arrugas
* Amarillamiento de los dientes
* Mal aliento
* Enfermedades de las encías
* Caries dental
* Pérdida dental
* Deterioro cognitivo
* Pérdida de memoria
* Confusión
* Depresión
* Ansiedad
* Insomnio
* Trastornos del sueño
* Fatiga
* Debilidad
* Calambres musculares
* Entumecimiento
* Hormigueo
* Dolor crónico
* Dolores de cabeza
* Migrañas
* Vértigo
* Tinnitus
* Pérdida de audición
* Problemas de visión
* Cataratas
* Glaucoma
* Degeneración macular
* Ojos secos
* Conjuntivitis
* Fatiga ocular
* Visión borrosa
* Mareos
* Problemas de equilibrio
* Problemas de coordinación
* Problemas de memoria
* Confusión
* Cambios de humor
* Irritabilidad
* Comportamiento agresivo
* Adicción
* Síntomas de abstinencia
* Antojos
* Recaída
* Aislamiento social
* Problemas financieros
* Problemas legales
* Conflictos familiares
* Pérdida de productividad
* Disminución del rendimiento
* Ausentismo
* Desempleo
* Pobreza
* Malnutrición
* Deshidratación
* Hipotermia
* Congelación
* Gangrena
* Amputación
* Muerte

ES

DESCARGO DE RESPONSABILIDAD
Por favor, lea este manual de usuario antes de usar el dispositivo. Si no sigue las instrucciones para la configuración, el uso y el cuidado adecuados del dispositivo, puede aumentar el riesgo de lesiones personales graves, muerte o daños materiales.

El fabricante no asume ninguna responsabilidad por los daños o lesiones causados por un manejo inadecuado o más allá del uso previsto.

CONFIGURACIÓN RECOMENDADA
Recomendamos el uso exclusivo de los cargadores, la batería y los componentes originales de Aspire. El uso de piezas de repuesto de terceros o no originales puede dañar el dispositivo, provocar sobrecalentamiento o condiciones de calor excesivo (167°C / 154°F).

Su dispositivo puede dañarse si se cae, se quema, se perfora, se aplasta o si entra en contacto con líquidos. Si sospecha que su dispositivo o su batería están dañados, deje de utilizarlos, ya que puede provocar un sobrecalentamiento o quemarse.

Para evitar que se produzcan fallos de encendido, siempre asegúrese de bloquear o apagar el dispositivo cuando no esté utilizándolo, especialmente cuando lo lleve en el bolsillo, el bolso o la cartera.

En caso de quemadura o deformación de una de las componentes del aparato, si los ve, por favor, deje de usarlos y contacte con el servicio de atención al cliente de Aspire.

SEGURIDAD EN EL MANTENIMIENTO DE LOS BOBINAS
Para no sufrir o quemar, quemar el algodón en una bobina que no funciona o está fría en el vacío. Deje reposar la cápsula durante 5 minutos después de llenarla para que el algodón pueda saturarse de e-líquido antes de usarlo.

SEGURIDAD DE LAS BATERÍAS REEMPLAZABLES
No use baterías no recargables. No use baterías dañadas o baterías con envoltura rota o dañada. Desconecte la batería del cargador cuando no esté cargando. No utilice su dispositivo de vapeo mientras se está cargando.

SEGURIDAD EN LA ELIMINACIÓN DEL PRODUCTO
No deseche este producto y su batería en el medio ambiente. Lleve desechos de los componentes del producto en puntos de recogida/eliminación específicos.

AVISO DE SALUD SEGURIDAD
No utilice el producto si está embarazada, en periodo de lactancia, si está generalmente enferma o si padece trastornos del sistema respiratorio y/o cardiovascular.

Si experimenta algún efecto secundario, deje de usar el producto. Si el efecto continúa después de dejar de usarlo, consulte a un profesional de la salud inmediatamente.

7. ADVERTENCIAS Y PRECAUCIONES SOBRE LA NICOTINA
Antes de utilizar el e-líquido que contiene nicotina, es aquello que debe tener en cuenta:
* Fumadores
* Prohibido a menores de edad.
* Nicotina es conocida por ser adictiva y se excreta en la leche materna.

8. EFECTOS DESCRITOS
* Aquellos que tienen problemas de salud en general, como problemas del corazón, pulmones, diabetes, ulceras o presión arterial alta, deben consultar a su médico antes de usar, ya que estas condiciones pueden ser exacerbadas por la nicotina.
* La nicotina puede afectar la circulación sanguínea y tener un efecto farmacológico en algunos lechos vasculares que pueden afectar la presión arterial.
* La nicotina puede afectar el sistema digestivo, incluyendo intestinos y vejiga.
* Aquellos que toman:
* Insulina, la nicotina puede afectar los niveles de azúcar en sangre.
* Rizamatan (para tratar el síndrome de las piernas inquietas)
* Triciclene (para tratar la esquizofrenia)

9. PROBLEMAS EFECTOS SECUNDARIOS
Si inhala demasiada nicotina o usa el producto demasiado, puede experimentar uno o más de los siguientes efectos:
* Mareos
* Náuseas (sensación de mareo)
* Dolor de cabeza
* Tos
* Irritación de la boca o la garganta
* Dolor de estómago
* Diarrea
* Hipo
* Palpitaciones
* Vomitar
* Arritmia en el pecho

10. PROBLEMAS EFECTOS SECUNDARIOS
Si se usa nicotina con regularidad, puede provocar:
* Náuseas, vomitar
* Dolor de cabeza
* Mareos
* Fatiga
* Temblores
* Arritmia cardíaca
* Dolor en el pecho
* Hipertensión
* Enfermedad arterial coronaria
* Accidente cerebrovascular
* Cáncer
* Congestión nasal
* Sinusitis
* Alergias
* Bronquitis
* Enfisema
* Trastornos gastrointestinales
* Úlcera péptica
* Gastritis
* Diarrea
* Estreñimiento
* Náuseas
* Vomitar
* Enfermedad por reflujo gastroesofágico (ERGE)
* Disfunción eréctil
* Eyaculación precoz
* Disminución de la libido
* Irregularidades menstruales
* Problemas de fertilidad
* Complicaciones del embarazo
* Bajo peso al nacer
* Parto prematuro
* Mortinato
* Síndrome de muerte súbita del lactante (SMSL)
* Impotencia
* Osteoporosis
* Debilidad muscular
* Dolor articular
* Artritis reumatoide
* Psoriasis
* Eczema
* Envejecimiento prematuro
* Arrugas
* Amarillamiento de los dientes
* Mal aliento
* Enfermedades de las encías
* Caries dental
* Pérdida dental
* Deterioro cognitivo
* Pérdida de memoria
* Confusión
* Depresión
* Ansiedad
* Insomnio
* Trastornos del sueño
* Fatiga
* Debilidad
* Calambres musculares
* Entumecimiento
* Hormigueo
* Dolor crónico
* Dolores de cabeza
* Migrañas
* Vértigo
* Tinnitus
* Pérdida de audición
* Problemas de visión
* Cataratas
* Glaucoma
* Degeneración macular
* Ojos secos
* Conjuntivitis
* Fatiga ocular
* Visión borrosa
* Mareos
* Problemas de equilibrio
* Problemas de coordinación
* Problemas de memoria
* Confusión
* Cambios de humor
* Irritabilidad
* Comportamiento agresivo
* Adicción
* Síntomas de abstinencia
* Antojos
* Recaída
* Aislamiento social
* Problemas financieros
* Problemas legales
* Conflictos familiares
* Pérdida de productividad
* Disminución del rendimiento
* Ausentismo
* Desempleo
* Pobreza
* Malnutrición
* Deshidratación
* Hipotermia
* Congelación
* Gangrena
* Amputación
* Muerte

NL

VELIGHEID DIMENSIES COIL
Wij raden ten sterkste aan om uit de verdamping van het batterij te verdrinken, maar wanneer u ze gebruikt, moet u ze pop-ta gas-sen. Laat de pompers ten minst 5 uit-1 stellen om ervoor te zorgen dat de wattes kunnen worden verzadigd met e-liquid.

VEILIGHEID VERVANGBARE BATTERIJEN
Gebruik geen niet-oplaadbare batterijen. Gebruik geen beschadigde batterijen of batterijen met een gescheurde of beschadigde verpakking. Ontkoppel de batterij van de lader wanneer hij niet is aangesloten. Gebruik uw vapeapparaat niet terwijl u aan het opladen bent.

VEILIGHEID VERWIJDERING PRODUCT
Gooi dit product en de batterij niet weg in het milieu. Breng de onderdelen van het product uitsluitend in specifieke inzameladressen.

VEILIGHEID GEZONDHEIDSWAARSCHUWING
Gebruik het product niet als